

Carnaby's Black-Cockatoo

Planting list

Feeding, Roosting and Nesting plants of Northam, Toodyay and York



Plant Name	Soil type	Height	Flower Time	Environment found in	Benefit to Carnaby's
<i>Acacia saligna</i>	Variety of soils, Clay, Sand	1.5-6m	Jul-Nov	Variety of habitats	Feeding
<i>Banksia attenuata</i>	Sand, sometimes over laterite	0.4-10m	Oct-Feb	Sand dunes, sand plains	Feeding
<i>Banksia dallanneyi</i>	Sand or sandy loam, laterite, granite, limestone, quartzite	0.3m	Apr-Sep	Hillslopes and breakaways	Feeding
<i>Banksia fraseri</i>	Sand, gravel, laterite, granite	0.6-6m	Apr-Sep	Hill slopes and breakaways	Feeding
<i>Banksia menziesii</i>	White, grey or yellow sand.	1.3-7m	Feb-Oct		Feeding
<i>Banksia prionotes</i>	Sand, loam, clay, laterite, granite	10m	Feb-Aug	Sandplains, sand dunes, undulating slopes.	Feeding
<i>Banksia sessilis</i>	Sand, limestone, laterite, granite	0.5-5m	Apr-Nov		Feeding
<i>Callistemon phoeniceus</i>	Loamy, sandy loam, clay loam	1-6m	Sep-Jan	Often along watercourses	Feeding
<i>Callitris pyramidalis</i>	Sand	8m			Feeding
<i>Corymbia calophylla</i>	Clay loam, sandy clay, gravel, sand over limestone, granite, laterite	40m	Mar-Dec	Flats, hills, slopes, breakaways, wetlands, fringing salt marches, beside drainage lines.	Feeding,
<i>Eremophila glabra</i>	Sand, clay, loam, limestone	0.1-3m	Mar-Dec	Winter-wet depressions, sandplains & dunes	Feeding
<i>Eucalyptus loxophleba</i>	Rocky loam, sand, sandy clay, laterite, dolerite, granite	15m	Jul-Feb	Near outcrops, flats, rises, slopes, hilltops, salt lakes, drainage lines	Feeding, Nesting
<i>Eucalyptus marginata</i>	Grey sand, clay or sandy loam.	40m	Jun-Jan	Hills, rises	Feeding, Roosting

Plant Name	Soil type	Height	Flower Time	Environment found in	Benefit to Carnaby's
<i>Eucalyptus rudis</i>	Sandy or loam soils.	5-20m	Jul-Sep	Wetter parts of south-western WA, flats, hillsides.	Roosting
<i>Eucalyptus salmonophloia</i>	Clay loam, clay, sand, gravel	4-30m	Aug-Dec	Plains & low hills.	Feeding, Nesting
<i>Eucalyptus salubris</i>	Clay loam, loam, sand, laterite	4-15m	Sep-Mar	Undulating plains, slopes.	Nesting
<i>Eucalyptus wandoo</i>	Sandy loam, clay loam, gravel, laterite, granite	3-25m	Dec-May	Stony rises, undulating terrain	Feeding, Nesting, Roosting
<i>Grevillea hookeriana</i>	Sand, sandy clay, laterite, gravel, loam	0.5-3m	Jul-Apr	Clay pans, moderate slopes, hills	Feeding
<i>Grevillea paniculata</i>	Sand, laterite, sandy clay, loam	0.5-3m	Jun-Nov	Sandplains, granite outcrops, hillsides.	Feeding
<i>Hakea circumalata</i>	Sand, sandy loam, sandy clay, gravel	0.6-1.5m	Aug-Sep		Feeding
<i>Hakea erinacea</i>	Sand, loam	0.2-1.6m	May-Nov	Laterite or granite outcrops	Feeding
<i>Hakea gilbertii</i>	Sand, laterite	0.6-2m	Aug-Sep		Feeding
<i>Hakea incrassata</i>	Sand, laterite, gravel, granite	0.35-1.5m	Jun-Nov		Feeding
<i>Hakea lissocarpa</i>	Sand, sandy loam, granite, laterite	0.4-1.5m	May-Sep		Feeding
<i>Hakea petiolaris</i>	Loam, granite	9m	Mar-Jul	Outcrops.	Feeding
<i>Hakea preissii</i>	Sand, loam, sandy clay	2-6m	Aug-Dec	Margins of salt marsh, plains.	Feeding
<i>Hakea prostrata</i>	Sandy soils, often over laterite, loam, gravel	1-3m	Jul-Oct	Hillslopes, granite outcrops, coastal dunes	Feeding
<i>Hakea scoparia</i>	Sand, gravel, loam, laterite, granite	1.2-3m	Apr-Sep	Breakaways, hills, outcrops, sandplains, road verges	Feeding
<i>Hakea trifurcata</i>	Sand over limestone or laterite, loam, gravel	1.5-3m	Apr-Oct		Feeding
<i>Xanthorrhoea preissii</i>	Sand, loam, gravelly sandy clay, laterite, granite	5m	Jun-Dec	Ranges, coastal plain, near watercourses	Feeding

Sources:

- <https://florabase.dpaw.wa.gov.au/>
- <https://www.dpaw.wa.gov.au/images/documents/plants-animals/threatened-species/carnabys/Plants used by Carnabys black cockatoo 20110415.pdf>



Australian Government

National
Landcare
Program



wheatbelt
natural resource
management